

Available online at www.sciencedirect.com



BEHAVIOURAL BRAIN RESEARCH

Behavioural Brain Research 167 (2006) 305-317

Research report

www.elsevier.com/locate/bbr

# Echolocation, vocal learning, auditory localization and the relative size of the avian auditory midbrain nucleus (MLd)

Andrew N. Iwaniuk<sup>a,\*</sup>, Dale H. Clayton<sup>b</sup>, Douglas R.W. Wylie<sup>a,c</sup>

<sup>a</sup> Department of Psychology, University of Alberta, Edmonton, Alta. T6G 2E9, Canada
<sup>b</sup> Department of Biology, University of Utah, Salt Lake City, UT, USA
<sup>c</sup> Centre for Neuroscience, University of Alberta, Edmonton, Alta. T6G 2E9, Canada

Received 20 July 2005; received in revised form 12 September 2005; accepted 19 September 2005 Available online 26 October 2005

#### Abstract

The avian nucleus mesencephalicus lateralis, pars dorsalis (MLd) is an auditory midbrain nucleus that plays a significant role in a variety of acoustically mediated behaviours. We tested whether MLd is hypertrophied in species with auditory specializations: owls, the vocal learners and echolocaters. Using both conventional and phylogenetically corrected statistics, we find that the echolocating species have a marginally enlarged MLd, but it does not differ significantly from auditory generalists, such as pigeons, raptors and chickens. Similarly, all of the vocal learners tend to have relatively small MLds. Finally, MLd is significantly larger in owls compared to all other birds regardless of how the size of MLd is scaled. This enlargement is far more marked in asymmetrically eared owls than symmetrically eared owls. Variation in MLd size therefore appears to be correlated with some auditory specializations, but not others. Whether an auditory specialist possesses a hypertrophied MLd appears to be depend upon their hearing range and sensitivity as well as the ability to resolve small azimuthal and elevational angles when determining the location of a sound. As a result, the only group to possess a significantly large MLd consistently across our analyses is the owls. Unlike other birds surveyed, owls have a battery of peripheral and other central auditory system specializations that correlate well with their hearing abilities. The lack of differences among the generalists, vocal learners and echolocaters therefore reflects an overall similarity in hearing abilities, despite the specific life history requirements of each specialization and species. This correlation between the size of a neural structure and the sensitivity of a perceptual domain parallels a similar pattern in mammals.

© 2005 Elsevier B.V. All rights reserved.

Keywords: Inferior colliculus; Prey capture; Echolocating; Parrot; Songbird; Hummingbird; Owls

## 1. Introduction

The evolution of sensory specializations is often correlated with an increase in size of the neural region(s) that mediate the specialization. Jerison [45] referred to this relationship as the 'principle of proper mass' whereby the size of a given neural structure is a reflection of the complexity of the behaviour that it subserves. Although there is considerable debate concerning how and why such a correlation exists [79], there are numerous examples of such correlations in vertebrates. For example, mammals with different somatosensory requirements can have dramatically different somatosensory representations, such as the expansion of the nose is the Star-nosed Mole (*Condylura cristata*) [11], the tail in Spider Monkeys (*Ateles* spp.) [67] and the forepaw in Raccoons (*Procyon lotor*) [68] This correlation between the size of a sensory region and behaviour is not restricted to the cortex or somatosensation, but is also present in visual [5], auditory [3,27] and gustatory systems [23]. Although there are numerous examples of this in mammals, systematic comparisons of behaviour and sensory regions are rare in birds [33]. Given that birds share many behavioural [55] and neural similarities [43,72] with mammals, it is likely that such correlations also exist in birds.

*Abbreviations:* IC, inferior colliculus; MLd, nucleus mesencephalicus lateralis, pars dorsalis; An, nucleus angularis; La, nucleus laminaris; MC, nucleus magnocellularis; SC, superior colliculus; SI, somatosensory cortex; TeO, tectum opticum

<sup>\*</sup> Corresponding author. Tel.: +1 780 492 7239; fax: +1 780 492 1768. *E-mail address:* brainsize@yahoo.ca (A.N. Iwaniuk).

<sup>0166-4328/\$ -</sup> see front matter © 2005 Elsevier B.V. All rights reserved. doi:10.1016/j.bbr.2005.09.015

Audition is of critical importance to a wide range of behaviours in birds, such as prey capture, individual and species recognition, vocal learning and mate selection. As a result of the range of behaviours that rely upon acoustic cues, studies of avian audition have revealed marked species differences in hearing range and the ability to localize sounds and discriminate pitch, intensity and temporal differences (see review in [21]). From this and other evidence, it is clear that several groups have evolved unique auditory specializations that are not present in other birds: localization of prey using only acoustic cues, vocal learning and echolocation. The first of these, the location of prey using acoustic cues, has been well documented in owls. Asymmetrically eared owls, like the Barn Owl (Tyto alba), can accurately acoustically locate prey up to 7 m away in complete darkness [63]. The vocal learners are all species that learn some, or all, of their vocal repertoire and includes the songbirds, parrots and hummingbirds. Specialized forebrain nuclei and their projections to and from auditory and motor structures in the thalamus, midbrain and hindbrain mediate the learning and production of vocalizations in all three taxa [7,25,31,42,44,80]. Unlike the owls and vocal learners, the echolocating birds have received relatively little attention. The only species known to echolocate are the Oilbird (Steatornis caripensis) [28] and the Swiftlets (Aerodramus and Collocalia spp.) [58,66,69]. All of these species primarily echolocate only when navigating through roosting sites and caves and do not appear to use echolocation to capture prey.

One region that is likely to be integral to all forms of acousticmediated behaviours is the nucleus mesencephalicus lateralis, pars dorsalis (MLd). MLd receives input from the two parallel auditory pathways [10,47] and as such plays a role in the integration of auditory information. MLd is homologous to both the mammalian inferior colliculus (IC) and the torus semicircularis (TS) in non-avian reptiles and amphibians [9,47,51]. Rylander [73] reported that MLd is hypertrophied in songbirds and Cobb [13,14] reported a similar hypertrophy of MLd in the echolocating Oilbird and Black-nest Swiftlet (Aerodramus maximus), which parallels the hypertrophy of IC in echolocating bats (Fig. 1). Lastly, the most detailed studies of the anatomy and function of MLd are in the Barn Owl, in which MLd plays a critical role in the auditory localization of prey (see reviews in [50,52]). Several authors reported that MLd appears to be enlarged in the Barn Owl and other asymmetrically eared owls, such as Saw-whet (Aegolius acadicus) and eared (Asio) owls, compared to symmetrically eared owls and other birds [13,84]. Thus, there is ample evidence to support a significant role of MLd in vocal learning, echolocation and prey localization that might result in its hypertrophy. A systematic analysis of size variation in MLd has not, however, been performed across species to test this hypothesis. Here, we present a detailed analysis of size variation in MLd across 84 species including the echolocating Swiftlets and Oilbird, all three vocal learning taxa and several owl species using both conventional and phylogenetically based statistics. Based on previous reports [13,14,73], we expected MLd to be significantly larger in all three auditory specialists compared to other birds (i.e., auditory 'generalists').



Fig. 1. Scatterplots of log-transformed inferior colliculus (IC) volume against (a) brain volume minus IC volume and (b) superior colliculus volume for echolocating (closed symbols) and non-echolocating (open circles) bats (data from [4]). It should be noted that all species of the suborder Megachiroptera are non-echolocating, with the exception of the Egyptian Fruit Bat (*Rousettus aegyptiacus*), which is indicated by the filled square.

#### 2. Materials and methods

#### 2.1. Specimens

We measured MLd and tectum opticum (TeO) in 72 specimens representing 54 species (Table 1). Representatives of all three auditory specializations were surveyed as well as representative species from other taxa in order to encompass as much interspecific variation in MLd size as possible. Two echolocating species were measured: Oilbird and Pygmy Swiftlet (Collocalia troglodytes) [66]. Closely related non-echolocating taxa were also measured. For the Oilbird, we measured several caprimulgiform species (Table 1), which are regarded by some as close relatives [8,17,56,77]. For the echolocating Swiftlet, we measured the closely related [65,66] and non-echolocating Glossy Swiftlet (C. esculenta) and the Common Swift (Apus apus), a swift that is not closely related to the Swiftlets and does not echolocate. Species from all three orders that exhibit vocal learning were also sampled: songbirds (11 spp.), parrots (15 spp.) and hummingbirds (5 spp.). Lastly, two asymmetrically eared owls, Barn Owl and Saw-whet Owl, and one symmetrically eared owl, the Southern Boobook Owl (Ninox boobook), were measured. Data for an additional 31 species were obtained from Boire [6] and Cobb [13], which included four vocal learners, one echolocating Swiftlet and several species classified as auditory generalists (see Table 1). For those species that were both measured and gleaned from the literature, the values presented are the means of the two data sets.

# Table 1

A list of the 84 species surveyed, the auditory category they each belong to, brain, tectum opticum (TeO) and nucleus mesencephalicus lateralis, pars dorsalis (MLd) volumes (all in mm<sup>3</sup>) and the ratio of MLd to TeO

Order	Species	п	Auditory category <sup>a</sup>	Brain volume	TeO volume	MLd volume	MLd:TeO	Source
Anseriformes								
Mallard	Anas platyrhynchos	2	G	6065	187.48	7.916	0.042	[6], This study
Plumed Whistling-duck	Dendrocygna eytoni	I	G	9421	164.81	7.850	0.047	This study
Apodiformss								
Black-nest Swiftlet	Aerodramus maximus	1	E	_	_	_	0.061	[13]
Common Swift	Apusapus	2	G	668	42.84	0.813	0.019	This study
Chimney Swift	Chaetura pelagica	2	G	343	30.47	0.668	0.033	[6,2]
Glossy Swiftlet	Collocalia esculenta	2	G	121	9.51	0.504	0.043	This study (FMNH, SEA132) [13]
Pygmy Swiftlet	C. troglodytes	2	E	139	9.71	0.543	0.056	This study (FMNH, SEA133, SEA134)
African Palm-swift	Cypsiurus parvus	1	G	300	-	-	0.052	[13]
Caprimulgiformes								
Feline Owlet-nightjar	Aegotheles insignis	1	G	1540	73.64	3.801	0.052	This study (BBM-NG 101365)
Indian Nightjar	Caprimulgus asiaticus	1	G	_	_	_	0.060	[13]
Nightjar	Caprimulgus sp.	1	G	734	58.81	2.319	0.039	[6]
Whip-poor-will	Caprimulgus vociferus	1	G	820	-	-	0.052	[2]
Spotted Nightjar	Eurostopodus argus	1	G	1013	60.97	3.515	0.058	This study
Common Poorwill	Phalaenoptilus nuttallii	1	G	—	-	_	0.057	[13]
Tawny Frogmouth	Podargus strigoides	2	G	5627	296.64	9.908	0.033	This study
Oilbird	Steatornis caripensis	2	E	3900	104.74	5.363	0.062	This study (USNM, 431365), [13]
Charadriiformes								
Least Sandpiper	Calidris minutilla	1	G	472	35.52	1.241	0.035	[6]
Killdeer	Charadrius vociferus	1	G	1073	100.92	1.817	0.018	[6]
Short-billed Dowitcher	Limnodromus griseus	2	G	1231	51.12	1.807	0.035	[6], this study
Common Tern	Sterna hirundo	1	G	1593	121.49	1.700	0.014	[6]
Masked Lapwing	Vanellus miles	I	G	2686	206.30	4.937	0.024	This study
Ciconiiformes								
Grey Heron	Ardea cinerea	1	G	8446	525.32	4.653	0.009	[6]
Cattle Egret	Bubulcus ibis	1	G	4025	213.76	2.068	0.010	This study
Columbiformes	Columba livia	r	G	2180	1/3 85	1 206	0.030	This study
rigeon	Columba livia	2	U	2109	143.65	4.290	0.050	[6]
Ringneck Dove	Streptopelia risoria	1	G	1141	95.11	2.319	0.024	[6]
Coraciiformes Laughing Kookaburra	Dacelo novaeguineae	3	G	4027	333.48	5.116	0.015	This study
Falconiformes								
Brown Goshawk	Accipiter fasciatus	1	G	4875	236.92	3.792	0.016	This study
Brown Falcon	Falco berigora	1	G	6007	387.05	7.950	0.021	This study
Australian Hobby Falcon	Falco longipennis	1	G	3035	200.36	3.230	0.016	This study
Galliformes								
Chukar	Alectoris chukar	1	G	2500	147.76	5.453	0.037	[6]
Ruffed Grouse	Bonasa umbellus	2	G	3136	182.33	8.574	0.047	This study
Golden Pheasant	Chrysolophus pictus	1	G	3369	211.64	9.664	0.046	[6]
Northern Bobwhite	Colinus virginianus	1	G	1091	81.13	3.826	0.047	[6]
Common Quail	Coturnix coturnix	1	G	811	67.90	2.650	0.039	[6]
Chicken	Gallus domesticus	1	G	2993	201.23	7.595	0.038	[6]
Turkey	Meleagns gallopavo	1	G	6097	552.93	19.766	0.036	[6]
Chase Charles	Numida meleagris	1	G	3951	233.86	10.239	0.044	[0] [6]
Chaco Chachalaca	Orians canicollis	1	U	33/4	203.07	0.430	0.032	[0]

# Table 1 (Continued)

Order	Species	п	Auditory category <sup>a</sup>	Brain volume	TeO volume	MLd volume	MLd:TeO	Source
Peafowl Ring-necked Pheasant	Pavo meleagris Phasianus colchicus	1 1	G G	7355 2762	284.95 129.83	15.141 8.164	0.053 0.063	[6] [6]
Gruiformes American Coot	Fulica americana	1	G	2719	127.65	5.454	0.043	This study
Passeriformes								
Brown Thornbill	Acanthiza pusilla	1	VL	434	34.81	0.515	0.015	This study
Eastern Spinebill	Acanthorhynchus tenuirostris	1	VL	489	29.46	0.822	0.028	This study
Blue-faced Honeyeater	Entomyzon cyanotis	1	VL	2227	96.99	3.391	0.035	This study
Eastern Yellow Robin	Eopsaltria australis	1	VL VI	839	39.40	1.758	0.045	This study
White plumed Honeyester	Gymnornina ilbicen	1		4017	202.27	5.261	0.016	This study
Noisy Miner	Manonna melanocephala	1	VL	2279	88 50	1 592	0.015	This study
Superb Lyrebird	Manonna metanocephata Menura novaehollandiae	1	VL VL	10163	384.66	4 107	0.013	This study
Spotted Paradalote	Pardalotus punctatus	1	VL.	401	17.09	0.837	0.049	This study
Grev Currawong	Strepera versicolor	1	VL	5425	270.86	4.200	0.016	This study
Double-barred Finch	Taenwpygia bichenovii	1	VL	409	25.81	0.655	0.025	This study
Zebra Finch	Taeniopygia guttata	1	VL	328	20.90	0.711	0.034	[6]
Pelecaniformes								
Double-crested Cormorant	Phalacrocorax auritus	1	G	7323	187.24	5.291	0.028	[6]
Procellariiformes Short-tailed Shearwater	Puffinus tenuirostris	1	G	4658	235.01	2.864	0.012	This study
Psittaciformes								
Masked Lovebird	Agapornis personata	1	VL	2824	82.57	2.575	0.031	This study
Australian King Parrot	Alisterus scapularis	1	VL	5305	200.90	4.455	0.022	This study
Blue-fronted Amazon	Amazona aestiva	1	VL	7955	273.47	4.142	0.015	This study
Galah	Cacatua roseicapilla	1	VL	6547	217.52	3.305	0.015	This study
Yellow-tailed Black-cockatoo	Calyptorhynchus funereus	1	VL	16111	309.66	5.340	0.017	This study
Eclectus Parrot	Eclectus roratus	1	VL	5796	208.59	4.492	0.022	This study
Musk Lorikeet	Glossopsitta concinna	2	VL	3198	99.52	2.669	0.027	This study
Budgerigar	Melopsittacus undulatus	1	VL	1220	43.53	2.386	0.055	[6]
Bourke's Parrot	Neopsephotus bourkii	1	VL	1213	56.42	1.884	0.033	This study
Cockatiel	Nymphicus hollandicus	3	VL	2263	75.31	2.111	0.028	This study
Blue-headed Parrot	Pionus menstruus	1	VL	5283	184.61	3.555	0.019	[6]
Crimson Rosella	Platycercus elegans	1	VL VI	3083	154.94	3.935	0.025	This study
Eastern Rosella	Platycercus eximius Polytolis guginsonii	2		2157	129.25	2.078	0.021	This study
Red rumped Parrot	Polylelis swainsonii Psephotus haematopotus	2	VL VI	1040	73 47	1.680	0.027	This study
African Grey Parrot	Psittacus erithacus	1	VL VI	6405	155.14	4 357	0.023	This study
Rainbow Lorikeet	Trichoglossus haematodus	2	VL	3728	123.42	2.578	0.020	This study
Rheiformes								·
Greater Rhea	Rhea americana	1	G	19228	953.28	38.190	0.040	[6]
Sphenisciformes			_					
Magellanic Penguin	Spheniscus magellanicus	1	G	16757	383.31	9.967	0.026	[6]
Strigiformes	4 11 11		0	22/5	<i>c</i> , , , , ,	14.007	0.001	
Saw-whet Owl	Aegolius acadicus	1	0	2343	64.49	14.227	0.221	This study
Southern Boobook Owl Barn Owl	Ninox boobook Tyto alba	2	0	5626 7143	148.15	9.261 26.150	0.063	This study This study
	1 yio uibu	1	0	/145	107.09	20.150	0.244	This study
Red-winged Tinamou	Rhynchotus rufescens	1	G	3377	327.43	9.964	0.030	[6]
Trochiliformes				100	10.00	0.000	0.005	
Anna's Hummingbird	Calypte anna	1	VL VI	183	12.33	0.329	0.027	This study
Blue-tailed Emerald	Chlorostilbon mellisugus	1		119	10.44	0.554	0.053	[0] This stude:
Green-ironted Lancebili	Doryjera ludoviciae	1	۷L	139	9.49	0.224	0.024	(FMNH 320498)
Buff-tailed Sicklebill	Eutoxeres condamini	2	VL	257	20.46	0.474	0.023	This study (FMNH 315304, 315300)

Table 1 (Continued)

Order	Species	n	Auditory category <sup>a</sup>	Brain volume	TeO volume	MLd volume	MLd:TeO	Source
Rufous-breasted Hermit	Glaucis hirsuta	1	VL	123	12.96	0.296	0.023	This study (USNM 616825)
Green-backed Firecrown	Sephanoides sephanoides	2	VL	134	10.19	0.238	0.023	This study (FMNH 316786, 316784)

The sources of the material are as follows—BBM: Bernice Bishop Museum (Honolulu, HI), USNM: National Museum of Natural History (Washington, DC), FMNH:Field Museum of Natural History (Chicago, IL) and references as indicated. Where more than one source is referenced, the values represent the mean. <sup>a</sup> Auditory categories are as follows—E: echolocaters, G: generalists, O: owls and VL: vocal learners.

Several specimens were loaned to us from the Field Museum of Natural History (Chicago, IL), National Museum of Natural History (Washington, DC) and the Bishop Museum (Honolulu, HI). All of the museum specimens were captured in the field, immersion fixed in 10% formalin for several days to several weeks and then stored in 70% ethanol for 3–45 years before we extracted the brains. Other fixed specimens were acquired from wildlife shelters, veterinary clinics and other labs. For all specimens, including the museum specimens, the brain was extracted, weighed to the nearest milligram and post-fixed in 4% paraformaldehyde in 0.1 M phosphate buffer (PB) for several days. The brains were then blocked in the sagittal plane, cryoprotected in 30% sucrose in PB, embedded in gelatin and sectioned in the coronal plane on a freezing stage microtome at a thickness of 40  $\mu$ m. Sections were collected in 0.1 M phosphate buffered saline, mounted onto gelatinized slides, stained with thionin and coverslipped with Permount.

#### 2.2. Measurements

Previous studies of MLd [14,73] compared the volume of MLd to the size of the optic lobe. The optic lobe in these studies is defined as the volume of the optic tectum and mesencephalon within those sections that the IC is located. The problem, however, is that this measure of the optic lobe is highly dependent upon the plane of section. For example, if the angle the sections are cut at is angled more rostro-ventrally in one specimen than another, the resulting measures of optic lobe volume will be markedly different. For this reason, we chose to scale the size of MLd against TeO. We chose TeO because studies in mammals have suggested that there is a trade-off between the size of the IC and the superior colliculus (SC) that reflects the degree of visual or auditory specialization (Fig. 1) and SC is homologous with the TeO [9]. Furthermore, Cobb [13] notes that TeO is a more meaningful measure because TeO is primarily visual whereas the optic lobe measurement encompasses parts of the mesencephalon that are unrelated to vision or hearing and provides a MLd: TeO ratio for eight species. Finally, the relative size of a given brain region can vary depending upon what is used as a scaling factor [18,38,64,79] and therefore the use of multiple scaling factors is recommended to assess how consistent the results are. We therefore compared the volume of MLd to both TeO and brain volume.

In accordance with previous studies [6] we defined TeO as the laminated portion of the optic lobe excluding the optic tract. The caudal and rostral poles of MLd were defined as those regions ventral to the third ventricle that had larger, darker and more densely packed cells than adjacent regions. The ventral and lateral borders were defined by the presence of a distinct lamina that forms a fibre bundle surrounding MLd [51] and the dorsal and lateral borders were defined by the tectal ventricle (Fig. 2). Although MLd can be subdivided into several subdivisions [51,85], the border between the central and external nuclei is faint [83] and we were unable to reliably distinguish the subdivisions in most of the specimens. Therefore, our measurements are restricted to the entire volume of MLd. For both TeO and MLd, our borders were similar to those used in previous allometric studies [6,13,14,73]. Whole brain volumes were measured by dividing the mass of each brain by the average density of fresh brain tissue [6,36,38,40].

Digital photographs were taken throughout the rostro-caudal extent of MLd and TeO for all specimens that we measured. Sampling intervals varied from every second to every fifth section depending on the size of the brain. Using this method, we measured 10–20 sections containing MLd and 30–40 sections containing TeO for each specimen. Measurements of each region were made using the NIH ImageJ freeware computer program. Volumes were calculated by multiplying the area by the sampling interval and section thickness. All of these procedures are similar to those used in previous studies [6,13,14,73].

#### 2.3. Statistical analysis

All species were classified according to four categories: owls, vocal learners, echolocaters and auditory generalists (i.e., species without any known auditory specializations). We then calculated MLd:TeO ratios (Table 1) and performed ANOVAs on these four categories to determine if there was a difference among the auditory specialists and other birds. To further examine relative size variation in MLd, we first logio-transformed MLd, TeO and brain minus MLd volumes (hereafter referred to as brain volume). Analyses of covariance (ANCOVAs) with auditory specialization and the scaling variables as covariates of MLd volume were then used to test for significant differences among groups. We also used ANOVAs of residuals derived from least-squares linear regression lines of MLd against TeO and brain volumes to assess differences among groups and Tukey–Kramer post hoc tests for pair-wise comparisons. In addition, we tested for differences among those orders represented by three or more species using ANOVAs of the MLd residuals to assess whether some orders tended to have larger MLds than others.



Fig. 2. A photomicrograph of the optic lobe of a songbird, the Spotted Pardalote (*Pardalotus punctatus*). The ventral and lateral borders of the nucleus mesencephalicus lateralis, pars dorsalis (MLd) are indicated by the solid line. Other structures shown are as follows: cerebellum (Cb), nucleus intercollicularis (ICo), nucleus isthmi, pars magnocellularis (Imc), nucleus isthmi, pars parvocellularis (Ipc), nucleus semiluminaris (SLu), stratum opticum (SOp), telencephalon (T), tectum opticum (TeO) and ventricle (V). Scale bar = 1 mm.

Because allometric analyses can be affected by phylogenetic relationships [32], we then repeated these analyses using phylogenetically based statistics. Phylogenetically corrected ANCOVAs and ANOVAs were performed using the PDAP software package (available from T. Garland). We performed these by simulating a null F distribution of traits along a phylogeny and using the critical values from this distribution as phylogenetically corrected critical Fs as opposed to critical Fs derived from conventional statistical tables [26]. This method has been used previously in comparative analyses of the brain and behaviour [36-38,64] and is less prone to type I error than conventional statistics when comparing data from many species [26]. To perform these analyses, a composite phylogeny was constructed using the inter-ordinal relationships in Sibley and Ahlquist [77] with resolution for some groups supplemented by additional references [1,2,12,48,66]. For each test, we constrained the simulation to biologically realistic values by setting the upper boundaries slightly higher and lower than that of the largest and smallest values respectively. The simulated F values were similar using both a gradual and a speciational model of evolutionary change, so we only report those of the gradual model.

# 3. Results

## 3.1. MLd:TeO ratio

The MLd:TeO ratio varied from 0.007 in the Grey Heron (Ardea cinerea) to 0.244 in the Barn Owl (Table 1). An ANOVA revealed a significant difference in MLd:TeO ratio among the auditory generalists and specialists (Table 2). Post hoc tests indicated that the owls have significantly larger MLd:TeO than any of the other groups and that the echolocaters have significantly larger ratios than the vocal learners (Fig. 3). Although the echolocating swifts (0.058) have a marginally higher ratio than the non-echolocating swifts (0.044), no significant difference was detected ( $t_{(4)} = -1.80$ , p = 0.15). The Oilbird, however, did have a significantly larger ratio than its putative relatives within the order Caprimulgiformes (single sample  $t_{(7)} = 3.12$ , p = 0.02). It should also be noted that there was substantial variability within the owls. Specifically, the asymmetrically eared Barn Owl and Saw-whet Owl have ratios three times that of the symmetrically eared Boobook Owl (Table 1).



Fig. 3. Boxplots of the nucleus mesencephalic lateralis, pars dorsalis (MLd) to tectum opticum (TeO) ratio for the echolocaters, owls, vocal learners and auditory generalists. The boxes represent the 95% confidence interval for each category, the horizontal line is the median and the filled circles are outliers. Significant differences were present between the owls and all other groups and between the echolocaters and vocal learners.

A significant difference in MLd:TeO was also found among orders ( $F_{(8,63)}$ =21.43, p<0.01). Post hoc pair-wise comparisons revealed that owls have significantly larger ratios than all other orders and the caprimulgiforms (which includes the Oilbird) have significantly larger ratios than the parrots (Psittaciformes). The significant difference among orders did not, however, exceed the phylogeny-corrected critical *F* (26.89). Overall then, owls tend to have larger MLd:TeO ratios than other groups, but this difference is minimal when phylogeny is taken into account.

#### 3.2. MLd and TeO

Comparing the size of MLd with TeO using an allometric approach yielded similar results to that of the MLd:TeO ratio. A scatterplot of MLd against TeO volume revealed that the two asymmetrically eared owls lie well above all of the other species sampled (Fig. 4a) and are, in fact, significant outliers. Although the Boobook Owl lies below the other two owls, it too has a larger MLd relative to TeO than the remaining species. An ANCOVA using auditory specializations and TeO volume as covariates of MLd volume did not yield a significant interaction



Fig. 4. A scatterplot of log-transformed nucleus mesencephalicus lateralis, pars dorsalis (MLd) against: (a) log-transformed tectum opticum (TeO) volume and (b) log-transformed brain volume minus MLd volume.

Table 2

The results of ANOVAs performed on relative nucleus mesencephalicus lateralis, pars dorsalis (MLd) volume using the scaling factors of: a ratio of MLd to tectum opticum (TeO) volume, residuals from a MLd volume and brain volume minus MLd volume regression and residuals from a MLd volume and TeO volume regression

Test	df	Calculated F	Critical F	Phylogeny-corrected critical F
MLd:TeO ratio	3, 81	56.51	2.72	9.76
MLd and TeO volume	3, 76	20.34	2.72	11.08
MLd and brain volume	3, 76	13.75	2.72	10.83

Critical Fs were derived from conventional statistical tables whereas the phylogeny-corrected critical Fs were calculated in PDANOVA (see text for details). Significant calculated Fs are shown in bold.

term, but there was a significant effect of auditory specialization (Table 3). This remained significant even when the calculated *F* was compared to the phylogeny-corrected critical *F* (Table 3). Thus, there is a significant difference in the relative size of MLd amongst the auditory specialists and generalists regardless of whether phylogenetic relationships are accounted for or not. In fact, this multiple regression model accounted for more variation ( $r^2 = 0.86$ ) than simply regressing MLd on TeO volume ( $F_{(1.78)} = 235.54$ , p < 0.01,  $r^2 = 0.75$ ).

An ANOVA of residuals derived from a least-squares linear regression of MLd on TeO volume also found a significant difference among the three auditory specialists and the generalists (Table 4, Fig. 5a). Post hoc Tukey–Kramers revealed that relative to TeO, MLd is significantly larger in owls compared to all other groups and is significantly smaller in vocal learners compared to generalists. No significant difference was detected between the echolocating Swiftlet and the other swifts (single sample  $t_{(2)} = -2.20$ , p = 0.16) or between the Oilbird and the caprimulgiforms (single sample  $t_{(4)} = -1.55$ , p = 0.20).

An ANOVA of the residuals also yielded a significant difference among orders ( $F_{(8,58)} = 15.89$ , p < 0.01). The pair-wise comparisons revealed that the owls have relatively larger MLd volumes than all other orders (Table 4). The order Caprimulgiformes also had significantly larger MLd residuals compared to the shorebirds, raptors and all vocal learners and the Galliformes had significantly larger MLd residuals than the hummingbirds. It should be noted, however, that the calculated *F* did not exceed the phylogeny-corrected critical *F* (34.08).

In summary, relative to TeO, MLd is significantly larger in owls than in other groups. MLd is not, however, relatively larger in echolocaters or either of the orders containing echolocating species (Apodiformes, Caprimulgiformes). MLd is also not significantly larger relative to TeO in the vocal learners.

### 3.3. MLd and brain volume

An examination of the allometric relationship between MLd and brain volume corroborated both analyses presented above. The asymmetrically eared Saw-whet and Barn Owls lie above all other species in a scatterplot of MLd against brain volume (Fig. 4b), although the difference between these owls and other birds is not as marked as the plot of MLd and TeO (Fig. 4a). An ANCOVA using brain volume and auditory specialization (owls, vocal learners and generalists) as covariates indicated no significant interaction term, but there was a significant effect of specialization (Table 3). Thus, there is a significant difference in relative MLd size amongst the auditory specialists and generalists. Furthermore, the  $r^2$  for a regression model including auditory specialization was higher (0.86) than a regression of simply MLd against brain volume ( $F_{(1,78)} = 248.72$ , p < 0.01,  $r^2 = 0.76$ ).

ANOVAs of the residuals from a least-squares linear regression line for all species also corroborated these findings (Table 2). A significant difference in relative MLd volume was found among the auditory specialists and generalists (Fig. 5b). Post hoc Tukey–Kramer tests revealed the owls have significantly larger MLd residuals than the generalists and vocal learners and vocal learners have significantly smaller MLd residuals than generalists. No significant difference in MLd residuals was detected between the echolocating Pygmy Swiftlet and the other swifts (single sample  $t_{(2)} = -1.46$ , p = 0.28). The Oilbird did, however, have a significantly larger MLd residual than the non-echolocating caprimulgiforms (single sample  $t_{(4)} = -5.98$ , p = 0.03).

When the residuals were grouped according to order, a significant difference was detected ( $F_{(8,58)} = 19.27, p < 0.01$ ). Pair-wise comparisons (Table 4) revealed that owls have larger relative MLd volumes than all other orders, except for Galliformes and Caprimulgiformes. The Galliformes have significantly larger

Table 3

The results of ANCOVAs performed on relative nucleus mesencephalicus lateralis, pars dorsalis (MLd) volume using the scaling factors of: a ratio of MLd to tectum opticum (TeO) volume), MLd and brain volume minus MLd volume and MLd volume and TeO volume

Test	Effect	df	Calculated F	Critical F	Phylogeny-corrected critical F
TeO	Interaction	3, 72	1.46	2.73	9.98
	Main effect	3, 75	<b>20.30</b>	2.73	10.39
Brain	Interaction	3, 72	0.27	2.73	12.23
	Main effect	3, 75	<b>21.40</b>	2.73	11.14

Critical Fs were derived from conventional statistical tables whereas the phylogeny-corrected critical Fs were calculated in PDANOVA (see text for details). Significant calculated Fs are shown in bold.

volume against both br	ain volume and tec	tum opticum (TeO) volu	Ime		-	)			
	Apodiformes	Caprimulgiformes	Charadriiformes	Falconiformes	Galliformes	Passeriformes	Psittaciformes	Strigiformes	Trochiliformes
Apodiformes		1		ı	I	-	-	+	
Caprimulgiformes	ı		+	+	ı	+	+	+	+
Charadriiformes		I		L	ı	I	ı	+	ı
Falconiformes	1	1	1		ı	1	ı	+	т
Galliformes	+	ı	+	+		+	-	+	+
Passeriformes	ı	+		ı	+		-	+	
Psittaciformes	I	+		L	+	I		+	ī
Strigiformes	+		+	+		+	+		+
Trochiliformes	ı	+	1	ı	+	L	1	+	
Those shown above the	shaded boxes are	the results from the MLo	I/TeO residuals and the	hose below the shae	ded boxes are from	n the MLd/brain re	siduals. For a list of	f the species within	i each order, refer to

Table 4

Table 1. Significant differences are indicated by a '+' whereas non-significant differences are indicated by a '-'.



Fig. 5. Boxplots of residuals from least-squares linear regressions of logtransformed nucleus mesencephalicus lateralis, pars dorsalis MLd volume against (a) log-transformed tectum opticum (TeO) volume, and (b) logtransformed brain volume minus MLd volume grouped according to: echolocaters, owls, vocal learners and auditory generalists. The boxes represent the 95% confidence interval for each category, the horizontal line is the median and the filled circles are outliers.

MLd residuals than the swifts, shorebirds (Charadriiformes), raptors (Falconiformes) and all three orders of vocal learners and the Caprimulgiformes have significantly larger MLd residuals than all three vocal learners. As with the previous analysis of MLd and TeO, this difference among orders was not supported by a comparison of the calculated F with the phylogeny-corrected critical F (31.13).

Based on this final analysis, it is clear that relative to brain volume, MLd is significantly larger in owls than in other birds. The Oilbird and its putative relatives, the caprimulgiforms, also have relatively large MLds, but only in comparison with the vocal learners. The other echolocaters, their relatives and all three groups of vocal learners do not, however, have relatively large MLds.

# 4. Discussion

Overall, our analyses show that compared to other birds, MLd is significantly larger in owls, but smaller in vocal learners. One of our analyses indicated that MLd is slightly larger in echolocaters compared to vocal learners and auditory generalists, but this was not corroborated by subsequent analyses. We therefore conclude that MLd is hypertrophied in owls, but not in vocal learners or echolocaters.

Although Cobb [13,14] and Rylander [73] reported that MLd was hypertrophied in both echolocaters and songbirds, there are several problems with their analyses (also see below). First, both authors used ratios to express the relative size of MLd. Ratios are problematic in allometric analyses because they are often correlated with the denominator and therefore do not reflect the true relative size of a structure [79]. Second, relatively few species were sampled in either study and therefore are unlikely to be representative of the variability in birds as a whole. For example, the relative size of MLd was not known for any of the vocal learning parrots and hummingbirds. Lastly, previous studies did not include any information on phylogenetic relatedness, which can have a significant effect on the analysis and interpretation of interspecific data [32]. Our analysis therefore represents a more comprehensive account of size variation in MLd among birds.

## 4.1. Vocal learning

Contrary to our prediction and the findings of Cobb [14] and Rylander [73], we found no evidence of MLd hypertrophy in songbirds, hummingbirds and parrots. In fact, vocal learners tend to have significantly smaller MLds relative to both brain and TeO volume than owls and auditory generalists. We contend that the reason for the discrepancy between our results and those of Cobb [14] and Rylander [73] is that their measurements were inaccurate. They calculated a ratio of MLd volume to that of the optic lobe, but only for those sections within which MLd was present. The danger of using such criteria is that the entire optic lobe was included. Furthermore, the area of their optic lobe measure is highly dependent upon the plane of section. As the plane of section moves from a coronal to a horizontal plane of section, the ratio increases because of the position of MLd within the optic lobe. Given that it is extremely difficult to obtain the same plane of section through all midbrain structures across species that vary in brain size and morphology, this is a highly unreliable measure. For this reason, we scaled MLd volume against our measures of the entire optic tectum and the entire brain volume.

and hummingbirds might reflect their basic auditory abilities. Although songbirds can localize sounds well [59,60], their spatial resolution is not as fine as that of owls and some other birds (Table 5). The audiogram of songbirds is not any broader than that of auditory generalists, despite being shifted towards slightly higher frequencies [21]. In addition, songbirds do not have particularly sensitive hearing in so far as the minimum audibility thresholds are quite high [21, Table 5]. They share these same characteristics not only with parrots (Table 5) and hummingbirds [70], but also many auditory generalists, such as galliforms, ducks and raptors (Table 5). Given that vocal learners have generally unremarkable hearing compared to auditory generalists, it is perhaps not surprising that MLd tends to be of average to below average size in vocal learners, despite the findings of previous studies [14,73]. In fact, the tuning properties of MLd neurons are not different to that reported in auditory generalists [88,89], unlike those of owls [82,83]. Although MLd must play some role in vocal learning because of its connections with the rest of the auditory system [47,81], it does not appear to be specialized for this role. This could reflect the prominent role that forebrain nuclei play in vocal learning, perception and discrimination (see review in [42]). In fact, the relatively large forebrains of both parrots and songbirds appear to result from the enlargement of multimodal regions, such as the caudo-medial nidopallium and caudo-medial mesopallium [36,38,39,71], both of which are involved in the learning and perception of vocalizations [42]. Volumetric analyses have demonstrated that both parrots and songbirds have particularly large forebrains and small TeO for their brain size [36,38,39]. Thus, vocal learners appear to have hypertrophied forebrain regions involved in auditory integration rather than MLd.

The relatively small size of MLd in songbirds, parrots

### 4.2. Echolocation

Cobb [13] reported that MLd was enlarged in the echolocating Black-nest Swiftlet and Oilbird compared to three swifts and three nightjars, respectively. Our study partially supports his findings. Using the MLd:TeO ratio, we found a significant difference between echolocaters and vocal learners and a significant difference between echolocating and non-echolocating species within the Caprimulgiformes, but not in the swifts. Although the

Group	Minimum absolute resolvable angle (°)	Higher frequency cutoff (kHz)	Auditory sensitivity at 500 Hz (dB)
Asymmetrically eared owls	1–6	12.5	<5
Symmetrically eared owl	7	7	<5
Songbirds	20-101	7–12	12–48
Budgerigar (Melopsittacus undulatus)	45–95	8	$\approx 18$
Oilbird (S. caripensis)	_	>8	≈35
Pigeon (Columba livid)	4–25	<6	23
Galliforms	$\approx 100$	6.5	32
Mallard (Anas platyrhynchos)	_	$\approx 6.5$	35
Raptors	2–14	7.5	25

The minimum absolute resolvable angle refers to the minimum angle difference that a bird can distinguish within a sound-attenuated room. The higher frequency cutoff is the frequency still detectable by the bird at 60 dB. The auditory sensitivity measure is the minimum dB detectable at a frequency of 500 Hz.

The auditory abilities of some of the taxa sampled (data from [20,48,52])

Oilbird did have a significantly larger MLd relative to brain volume than the caprimulgiforms, it should be noted that whether the Oilbird truly is a caprimulgiform or not is questionable [56]. From these mixed results and the fact that there were no significant differences detected between echolocating birds and auditory generalists, we conclude that echolocating birds may have slightly enlarged MLds compared to their relatives, but this enlargement does not differ in magnitude from that of other birds that do not echolocate. Therefore, unlike most bats, echolocating birds do not have a significantly enlarged MLd/IC, which likely reflects differences in the structure and function of echolocating vocalizations between bats and birds.

In both Oilbirds and Swiftlets, the echolocating calls are noisy broad-band clicks [15,22,24,53,66] and these calls are not used for the detection and capture of prey. Oilbirds feed primarily on fruit [19], which does not require echolocation to find, and Swiftlets likely use a combination of visual (e.g., motion parallax) and tactile (e.g., somatosensation by rictal bristles) to locate and capture insects in the air. The calls are primarily used when entering and exiting roosting sites within caves and likely function as an aid in navigating through dimly lit caves and around conspecifics [57]. This is further supported by experimental studies on the sensitivity and acuity of echolocation in both Oilbirds and Swiftlets. Swiftlets are able to successfully negotiate flight paths through large wooden dowels (6–10 mm), but their performance drops off significantly for thinner dowels [16,22,29,57,78]. Similarly, Oilbirds can avoid large objects in their flight path using echolocation, but not smaller objects [53].

In contrast to birds, the echolocation system of most bats is far more sensitive. Similar experiments to those conducted on Swiftlets have shown that bats are able to detect dowels/wires as thin as 0.06 mm [61]. In addition, most bat species do not use broad-band echolocating calls; instead, bats produce calls of a constant frequency (cf), frequency sweeps (fm) and a combination of constant frequency and frequency modulation (cf-fm) [61]. One notable exception to this is echolocating in fruit bats of the genus Rousettus, which employ broad-band clicks [35] when entering and exiting cave roosts and not for prey capture. Rousettus bats are therefore comparable to both the Swiftlets and the Oilbird in terms of the structure and function of their echolocating calls. These fruit bats also cannot resolve azimuthal angles as accurately as other echolocating bats [34], lack the spectral acuity of other echolocating bats [30] and do not have an enlarged IC (Fig. 1). Currently we do not have information on minimum resolvable angles of echolocating birds, which would provide further insight into how comparable echolocating birds are with Rousettus.

It must also be emphasized that our mixed results do not indicate that there are no neural specializations for echolocation in birds. Given the broad-band structure of both Swiftlet [15,22,24,66] and Oilbird calls [53], they may rely upon interaural timing differences (ITDs) more than interaural intensity differences (IIDs) to navigate. In birds, both ITDs and IID s are processed by MLd, but only after they are received by the cochlear nuclei, nucleus laminaris (La) and nucleus magnocellularis (MC) (see review in [52]). The Oilbird does have significantly larger La and MC than most other birds, the exceptions being the asymmetrically eared owls [54,86] and a similar hypertrophy of homologous cochlear nuclei also occurs in echolocating bats, but not in non-echolocating bats [3]. It is therefore likely that the echolocating Swiftlets also possess relatively large La and MC, which would enable them to respond quickly to ITDs in echoing calls without necessarily requiring a large MLd.

#### 4.3. Auditory localization

Of all the species sampled, the owls were the only group to have a significantly larger MLd, regardless of the statistical method used. As predicted, this likely reflects the use of acoustic cues to locate prey. Owls can resolve smaller azimuthal angles than other birds [49, Table 5], possess a detailed map of auditory space in MLd [82] and have the broadest and most sensitive hearing of any group of birds [21], Table 5. In addition, all owls possess specializations in the peripheral auditory system, such as unique malleus morphology [75], a long cochlea [74,76] and a relatively large tympanic membrane [74,75]. Given the precision of sound localization in owls [63] and this assortment of auditory specializations, it is not surprising that MLd is significantly enlarged. This therefore represents a third case of central nervous system structure hypertrophy in owls, the other two being the Wulst [39,84] and the cochlear nuclei La and MC [54,86].

Not all owls share the same precision and unique adaptations of the Barn and Saw-whet Owls and this appears to relate to the hypertrophy of both MLd and the cochlear nuclei. Asymmetrically eared owls, such as the Barn Owl, Saw-whet Owl, Great Grey Owl (Strix nebulosa) and Long-eared Owl (Asio otus), not only have asymmetrical ear openings [62], but also the longest cochleae, basilar membranes and papillae [74]. Behaviourally, all of these species can hunt in complete darkness [63] and can discern angular differences at a fine level (Table 5). Neurons within MLd of these species encode for both azimuth and elevation [82,87]. In contrast, the symmetrically eared owls, such as the Boobook Owl, Screech Owl (Otus asio), Burrowing Owl (Athene cunicularia) and Great-horned Owl (Bubo virginianus), do not have markedly asymmetrical ears [62]. Although a detailed description of their inner ear anatomy is wanting, the symmetrically eared owls only have moderately enlarged cochleae [74]. These species also tend to be more diurnal in their habits, are unable to capture prey in complete darkness [63] and cannot resolve azimuthal angles as finely as the asymmetrically eared owls (Table 5). Electrophysiological studies in the Burrowing Owl and Great-horned Owl also failed to find neurons that were tuned to elevation [81,82]. In this study, we also found that the hypertrophy of MLd was far less in the Boobook Owl than it was in the Barn and Saw-whet Owls relative to both TeO and brain volume. Comparisons of cochlear nuclei found similar variation; all of the symmetrically eared owls had smaller nuclei than the asymmetrically eared owls [54,86]. Thus, it appears that the symmetrically eared owls do not share the same spectrum of peripheral and central specializations for auditory localization as the asymmetrically eared owls.

The difference between asymmetrically and symmetrically eared owls highlights the importance of MLd function in its hypertrophy. In owls, IIDs and ITDs are combined to construct a map of auditory space within MLd [50,82,83]. As discussed above, the asymmetrically eared owls appear to use a more precise and detailed auditory map and therefore require a relatively large MLd. In other birds, such as vocal learners, auditory integration appears to take place in the forebrain rather than MLd. Thus, there may be a functional difference in the computations that MLd is performing among different taxa that underlies the variation in relative size. Further research into MLd tuning and function is, however, required amongst a range of avian species to support this hypothesis.

## 5. Conclusions

Despite the fact that two of our predictions were not supported by the data, the results do provide a better understanding of the function of MLd is acoustically mediated behaviours. Specifically, it appears that specialized tuning and hypertrophy of MLd is only present in asymmetrically eared owls and this is directly correlated with their ability to precisely locate prey using only acoustic cues. In doing so, our study supports a broader pattern observed in the evolution of behaviour and sensory systems in mammals and other vertebrates; the enlargement of a sensory region is correlated with the sensitivity of that sensory modality. As mentioned previously, Jerison [45] first suggested that the enlargement of brain regions is correlated with behavioural complexity and several studies have supported this principle. However, this is not always the case. For instance, the large representation of the forepaw in the primary somatosensory cortex (SI) of Raccoons was once considered to be indicative of their digital dexterity [68]. Thus, an expansion of the forepaw representation is correlated with an increase in the complexity of motor output such that Raccoons should be capable of similar digit movements as primates. A behavioural analysis of forepaw use in Raccoons [41], however, found that they are not as dexterous as many other species with smaller forepaw representations in SI, such as Rats (Rattus norvegicus). The Raccoon does have acute tactile discrimination abilities [46], but their motor output is not that dissimilar from many other mammals. Thus, the forepaw representation in Raccoons appears to reflect their ability to detect subtle tactile differences and not dexterity. This is the most likely functional outcome of the principle of proper mass [45]; the enlargement of a sensory region is correlated with sensitivity and not necessarily with any additional motor capacity and our analysis of MLd variation in birds supports this. The systematic sampling of sensory systems across birds will therefore provide significant insight into the correlated evolution of the brain and behaviour.

#### Acknowledgements

We wish to thank the curatorial staff of the Bishop Museum, Field Museum of Natural History and National Museum of Natural History for loaning us specimens, Healesville Sanctuary and David Madill for collecting many specimens on our behalf and Doug Altshuler, Barrie Frost and Gerard Gory for sending us heads of *Calypte anna*, *Aegolius acadicus* and *Apus apus*, respectively, and two anonymous reviewers for their helpful comments on a previous version of this manuscript. This research was funded by fellowships and grants from the Alberta Ingenuity Fund and the Natural Sciences and Engineering Research Council of Canada (NSERC) to AM, an NSF CAREER award to DHC and grants from NSERC and the Canada Research Chairs Program to DRWW.

## References

- Altshuler DL, Dudley R, McGuire JA. Resolution of a paradox: hummingbird flight at high elevation does not come without a cost. Proc Natl Acad Sci USA 2004;101:17731–6.
- [2] Barker FK, Cibois A, Schikler P, Feinstein J, Cracraft J. Phylogeny and diversification of the largest avian radiation. Proc Natl Acad Sci USA 2004;101:11040–5.
- [3] Baron G. Differential phylogenetic development of the acoustic nuclei among Chiroptera. Brain Behav Evol 1974;9:7–40.
- [4] Baron G, Stephan H, Frahm HD. Comparative Neurobiology in Chiroptera. Basel: Birkhäuser Verlag; 1996, 1596 pp.
- [5] Barton RA. Visual specialization and brain evolution in primates. Proc R Soc London B 1998;265:1933–7.
- [6] Boire D. Comparaison quantitative de l'encephale de ses grades subdivisions et de relais visuals, trijumaux et acoustiques chez 28 especes. PhD Thesis. Montreal: Universite de Montreal; 1989.
- [7] Brauth SE, Liang W, Roberts TF. Projections of the oval nucleus of the hyperstriatum ventrale in the budgerigar: relationships with the auditory system. J Comp Neurol 2001;432:481–511.
- [8] Brumfield RT, Swofford DL, Braun MJ. Evolutionary relationships among the potoos (Nyctibiidae) based on isozymes. Ornithol Monogr 1997;48:129–45.
- [9] Butler AB, Hodos W. Comparative Vertebrate Neuroanatomy. New York: Wiley Press; 1996, 514 pp.
- [10] Carr CE, Code RA. The central auditory system of reptiles and birds. In: Dooling RJ, Fay RR, Popper AN, editors. Comparative Hearing in Birds and Reptiles. New York: Springer-Verlag; 2000. p. 197– 248.
- [11] Catania K, Kaas JH. Organization of the somatosensory cortex in the star-nosed mole. J Comp Neurol 1995;351:549–67.
- [12] Christidis L, Schodde R, Shaw DD, Maynes SF. Relationships among the Australo-Papuan parrots, lorikeets and cockatoos (Aves:Psittaciformes). Condor 1991;93:302–17.
- [13] Cobb S. On the size of the auditory nuclei in some Apodiformes and Caprimulgiformes. Auk 1968;85:132–3.
- [14] Cobb S. A comparison of the size of an auditory nucleus (*n. mesen-cephalicus lateralis, pars dorsalis*) with the size of the optic lobe in twenty-seven species of birds. J Comp Neurol 1964;122:271–9.
- [15] Coles RB, Konishi M, Pettigrew JD. Hearing and echolocation in the Australian grey swiftlet, *Collocalia spodiopygia*. J Exp Biol 1987;129:365–71.
- [16] Collins CT, Murphy R. Echolocation acuity of the Palawan swiftlet (Aerodramus palawanensis). Avocetta 1994;17:157–62.
- [17] Cracraft J. Towards a phylogenetic classification of birds of the world (class Aves). Auk 1981;98:681–714.
- [18] Deaner RO, Nunn CL, van Schaik CP. Comparative tests of primate cognition: different scaling methods produce different results. Brain Behav Evol 2000;55:44–52.
- [19] del Hoyo J, Elliott A, Sargatal J. Handbook of the Birds of the World, vol. 5: Barn-owls to Hummingbirds. Barcelona: Lynx Edicions; 1999, 762 pp.
- [20] Dooling RJ. Auditory perception in birds. In: Kroodsma DE, Miller EH, Ouellet H, editors. Acoustic communication in birds, vol. 1. New York: Academic Press; 1982. p. 95–130.
- [21] Dooling RJ, Lohr B, Dent ML. Hearing in birds and reptiles. In: Dooling RJ, Fay RR, Popper AN, editors. Comparative hearing in birds and reptiles. New York: Springer-Verlag; 2000. p. 308–59.

- [22] Fenton MB. Acuity of echolocation in *Collocalia hirundinacea* (Aves: Apodidae), with comments on the distributions of echolocating swiftlets and molossid bats. Biotropica 1975;7:1–7.
- [23] Finger TE. Feeding patterns and brain evolution in ostariophysean fishes. Acta Physiol Scand Suppl 1975;638:59–66.
- [24] Fullard JH, Barclay RMR, Thomas DW. Echolocation in free-flying Atiu swiftlets (*Aerodramus sawtelli*). Biotropica 1993;25:334–9.
- [25] Gahr M. Neural song control system of hummingbirds: comparison to swifts, vocal learning (songbirds) and nonlearning (suboscines) passerines, and vocal learning (budgerigars) and nonlearning (dove, owl, gull, quail, chicken) nonpasserines. J Comp Neurol 2000;426:182– 96.
- [26] Garland Jr T, Dickerman AW, Janis CM, Jones JA. Phylogenetic analysis of covariance by computer simulation. Syst Biol 1993;42:265– 92.
- [27] Glendenning KK, Masterton RB. Comparative morphometry of mammalian central auditory systems: variation in nuclei and form of the ascending system. Brain Behav Evol 1998;51:59–89.
- [28] Griffin DR. Acoustic orientation in the oil bird, *Steatornis*. Proc Natl Acad Sci USA 1953;39:884–93.
- [29] Griffin DR, Suthers RA. Sensitivity of echolocation in cave swiftlets. Biol Bull 1970;139:495–501.
- [30] Griffin DR, Novick A, Kornfield M. The sensitivity of echolocation in the fruit bat, *Rousettus*. Biol Bull 1958;115:107–13.
- [31] Hall WS, Cohen PL, Brauth SE. Auditory projections to the anterior telencephalon in the budgerigar (*Melopsittacus undulatus*). Brain Behav Evol 1993;41:97–116.
- [32] Harvey PH, Pagel MD. The comparative method in evolutionary biology. Oxford: Oxford University Press; 1991, 239 pp.
- [33] Healy S, Guilford T. Olfactory-bulb size and nocturnality in birds. Evolution 1990;44:339–46.
- [34] Heffner RS, Koay G, Heffner HE. Sound localization in an Old-World fruit bat (*Rousettus aegyptiacus*): acuity, use of binaural cues, and relationship to vision. J Comp Psychol 1999;113:297–306.
- [35] Holland RA, Waters DA, Rayner JM. Echolocation signal structure in the Megachiropteran bat *Rousettus aegyptiacus* Geoffroy 1810. J Exp Biol 2004;207:4361–9.
- [36] Iwaniuk AN. The evolution of brain size and structure in birds. PhD Thesis. Clayton: Monash University; 2003.
- [37] Iwaniuk AN, Arnold KE. Is cooperative breeding associated with bigger brains? A comparative test in the Corvida (Passeriformes). Ethology 2004;110:203–20.
- [38] Iwaniuk AN, Dean KM, Nelson JE. Interspecific allometry of the brain and brain regions in parrots (Psittaciformes): comparisons with other birds and primates. Brain Behav Evol 2005;65:40–59.
- [39] Iwaniuk AN, Hurd PL. The evolution of cerebrotypes in birds. Brain Behav Evol 2005;65:215-30.
- [40] Iwaniuk AN, Nelson JE. Can endocranial volume be used as an estimate of brain size in birds? Can J Zool 2002;80:16–23.
- [41] Iwaniuk AN, Whishaw IQ. How skilled are the skilled limb movements of the raccoon (*Procyon lotor*). Behav Brain Res 1999;99:35–44.
- [42] Jarvis ED. Brains and birdsong. In: Marler P, Slabbekoorn H, editors. Nature's music: the science of birdsong. San Diego: Elsevier Press; 2004. p. 226–71.
- [43] Jarvis ED, Güntürkün O, Bruce L, Csillag A, Karten H, Kuenzel W, et al. Avian brains and a new understanding of vertebrate brain evolution. Nat Rev Neurosci 2005;6:151–9.
- [44] Jarvis ED, Ribeiro S, da Silva ML, Ventura D, Vielliard J, Mello CV. Behaviourally driven gene expression reveals song nuclei in hummingbird brain. Nature 2000;406:628–32.
- [45] Jerison HJ. Evolution of the brain and intelligence in vertebrates. New York: Academic Press; 1973, 482 pp.
- [46] Johnson Jr JI, Michels KM. Discrimination of small intervals and objects by raccoons. Anim Behav 1958;6:164–70.
- [47] Karten HJ. The organization of the ascending auditory pathway in the pigeon (*Columba livid*). I. Diencephalic projections of the inferior colliculus (nucleus mesencephali lateralis, pars dorsalis). Brain Res 1967;6:409–27.

- [48] Kimball RT, Braun EL, Zwartjes PW, Crowe TM, Ligon JD. A molecular phylogeny of the pheasants and partridges suggests that these lineages are not monophyletic. Mol Phylogenet Evol 1999;11:38–54.
- [49] Klump GM. Sound localization in birds. In: Dooling RJ, Fay RR, Popper AN, editors. Comparative hearing in birds and reptiles. New York: Springer-Verlag; 2000. p. 249–307.
- [50] Knudsen EI. Instructed learning in the auditory localization pathway of the barn owl. Nature 2002;417:322–8.
- [51] Knudsen EI. Subdivisions of the inferior colliculus in the barn owl (*Tyto alba*). J Comp Neurol 1983;218:174–86.
- [52] Konishi M. Coding of auditory space. Annu Rev Neurosci 2003;26:31–55.
- [53] Konishi M, Knudsen EI. The oilbird: hearing and echolocation. Science 1979;204:425–7.
- [54] Kubke MF, Massoglia DP, Carr CE. Bigger brains or bigger nuclei? Regulating the size of auditory structures in birds. Brain Behav Evol 2004;63:169–80.
- [55] Marler P. Social cognition: are primates smarter than birds? Curr Ornithol 1996;13:1–32.
- [56] Mayr G, Manegold A, Johansson US. Monophyletic groups within 'higher land birds'—comparison of morphological and molecular data. J Zool Syst Evol Res 2003;41:233–48.
- [57] Lord M. The function of echonavigation among swiftlets. Anim Behav 1967;15:416–20.
- [58] Lord M, Pye JD. Echolocation and the systematics of swiftlets. In: Stonehouse B, Perrins C, editors. Evolutionary ecology. Baltimore: University Park Press; 1977. p. 225–38 (Chapter 19).
- [59] Nelson BS, Stoddard PK. Accuracy of auditory distance and azimuth perception by a passerine bird in natural habitat. Anim Behav 1998;56:467–77.
- [60] Nelson BS, Suthers RA. Sound localization in a small passerine bird: discrimination of azimuth as a function of head orientation and sound frequency. J Exp Biol 2004;207:4121–33.
- [61] Neuweiler G. The biology of bats. Oxford: Oxford University Press; 2000, 320 pp.
- [62] Norberg RÅ. Occurrence and independent evolution of bilateral ear asymmetry in owls and implications on owl taxonomy. Phil Trans R Soc London B 1977;280:375–408.
- [63] Payne RS. Acoustic location of prey by barn owls (*Tyto alba*). J Exp Biol 1971;54:535–73.
- [64] Pellis SM, Iwaniuk AN. Brain system size and adult-adult play in primates: a comparative analysis of the non-visual neocortex and the amygdala. Behav Brain Res 2002;134:31–9.
- [65] Price JJ, Johnson KP, Bush SE, Clayton DH. Phylogenetic relationships of the Papuan Swiftlet and implications for the evolution of avian echolocation. Ibis 2005;1471:790–6.
- [66] Price JJ, Johnson KP, Clayton DH. The evolution of echolocation in swiftlets. J Avian Biol 2004;35:135–43.
- [67] Pubols Jr BH, Pubols LM. Neural organization of somatic sensory representation in the spider monkey. Brain Behav Evol 1972;5:342– 66.
- [68] Pubols Jr BH, Welker WI, Johnson Jr JI. Somatic sensory representation of forelimb in dorsal root fibers of raccoon, coatimundi, and cat. J Neurophysiol 1965;28:312–41.
- [69] Pye JD. Echolocation signals and echoes in air. In: Busnel RG, Fish JF, editors. Animal sonar systems. New York: Plenum Press; 1979. p. 309–53.
- [70] Pytte CL, Ficken MS, Moiseff A. Ultrasonic singing by the blue-throated hummingbird: a comparison between production and perception. J Comp Physiol A 2004;190:665–73.
- [71] Rehkämper G, Frahm HD, Mann MD. Evolutionary constraints of large telencephala. In: Roth G, Wullimann MF, editors. Brain evolution and cognition. New York: Wiley; 2001. p. 265–93.
- [72] Reiner A, Perkel DJ, Bruce LL, Butler AB, Csillag A, Kuenzel W, et al. Revised nomenclature for avian telencephalon and some related brainstem nuclei. J Comp Neurol 2004;473:377–414.
- [73] Rylander MK. The nucleus mesencephalicus lateralis, pars dorsalis in neotropical passerines. Brain Behav Evol 1979;16:315–8.

- [74] Schwartzkopf J. Structure and function of the ear and of the auditory brain areas in birds. In: DeReuck AVS, Knight J, editors. Hearing mechanisms in vertebrates. London: J&A Churchill Ltd.; 1968. p. 41–63.
- [75] Schwartzkopf J. On the hearing of birds. Auk 1955;72:340-7.
- [76] Schwartzkopf J, Winter P. Zur Anatomie der Vogel-Cochlea unter natürlichen Bedgingungen. Biol Zentral-bl 1960;79:607–25.
- [77] Sibley CG, Ahlquist JE. Phylogeny and classification of birds. New Haven: Yale University Press; 1990, 1080 pp.
- [78] Smyth DM, Roberts JR. The sensitivity of echolocation by the grey swiftlet Aerodramus spodiopygius. Ibis 1983;125:339–45.
- [79] Striedter GF. Principles of brain evolution. Sunderland: Sinauer Associates; 2004, 436 pp.
- [80] Striedter GF. The vocal control pathways in budgerigars differ from those in songbirds. J Comp Neurol 1994;343:35–56.
- [81] Vates GE, Broome BM, Mello CV, Nottebohm F. Auditory pathways of caudal telencephalon and their relation to the song system of adult male zebra finches. J Comp Neurol 1996;366:613–42.
- [82] Volman SF, Konishi M. Comparative physiology of sound localization in four species of owls. Brain Behav Evol 1990;36:196–215.

- [83] Volman SF, Konishi M. Spatial selectivity and binaural responses in the inferiorcolliculus of the great horned owl. J Neurosci 1989;9:3083–96.
- [84] Wagner H, Luksch H. Effect of ecological pressures on brains: examples from avian neuroethology and general meanings. Z Naturforsch C 1998;53:560–81.
- [85] Wagner H, Güntürkün O, Nieder B. Anatomical markers for the subdivisions of the barn owl's inferior–collicular complex and adjacent periand subventricular structures. J Comp Neurol 2003;465:145–59.
- [86] Winter P. Vergleichende Qualitative und quantitative untersuchungen an der hörbahn von vögeln. Z Morph Ökol Tiere 1963;52:365–400.
- [87] Wise LZ, Frost BJ, Shaver SW. The representation of sound frequency and space in the mid brain of the Saw-whet Owl. Soc Neurosci Abst 1988;14:1095.
- [88] Woolley SMN, Casseday JH. Response properties of single neurons in the zebra finch auditory midbrain: response patterns, frequency coding, intensity coding, and spike latencies. J Neurophysiol 2004;91:136–51.
- [89] Woolley SMN, Casseday JH. Processing of modulated sounds in the zebra finch auditory midbrain: responses to noise, frequency sweeps and sinusoidal amplitude modulations. J Neurophysiol 2005;94:1143–57.